

Note

Injective Choice Functions

KARSTEN STEFFENS*

Institut für Mathematik, Technische Universität Hannover, Hannover, West Germany

Communicated by the Managing Editors

Received September 13, 1972

The present paper is concerned with a combinatorial question called the "marriage problem." A criterion will be proved for the existence of an injective choice function of families with at most finitely many infinite members and a generalization of a theorem of H. A. Jung and R. Rado. We give a new proof of a theorem of J. Folkman.

Let $F = (F(i) : i \in I)$ be a family of nonempty sets $F(i)$ with the index set I . We call a function f a *choice function* of F if the domain of f is I and $f(i)$ is an element of $F(i)$, for all elements i of I . Since there is no injective choice function of the family $F = \{(0, \{0, 1\}), (1, \{0\}), (2, \{1\})\}$, we are interested in a criterion (necessary and sufficient) for the existence of an injective choice function of a given family. We shall use the letters F, G, H for families of nonempty sets, the letters I, J, K for index sets, and the letters f, g, h for choice functions. We write $\text{rng } f$ for the range of f , $\text{dom } f$ for the domain of f , and $f \upharpoonright K$ ($F \upharpoonright K$) for the restriction of f (F) to the set K . $P(I)$ denotes the power set of I , $P_\omega(I)$ the set of all finite subsets of I , and $\omega = \{0, 1, 2, 3, \dots\}$, the set of all natural numbers. The letter n always denotes a natural number. $IA(F, f)$ means that f is an injective choice function of the family F . Let us call a family F *critical* if (1) $\exists f IA(F, f)$ and (2) $\neg \exists f (IA(F, f) \ \& \ \text{rng } f \subsetneq \bigcup \text{rng } F)$.

LEMMA 1. *Let $F = (F(i) : i \in I)$ be a family. Then*

$$\exists f IA(F, f) \Rightarrow \neg \exists K \subseteq I \exists i \in I \setminus K \left(F(i) \subseteq \bigcup \text{rng } F \upharpoonright K \ \& \ F \upharpoonright K \text{ critical} \right).$$

The proof is straightforward. We hope that the necessary condition of Lemma 1 is also sufficient. First we attack the problem for finite families.

* This note is a part of the author's doctoral thesis. The author wishes to express his gratitude to Prof. Dr. Walter Oberschelp as his thesis advisor who has directed the author's attention to combinatorial problems.

THEOREM 1 (P. Hall). *Let $F = (F(i) : i \in n)$ be a finite family. Then*

$$\exists f \text{ IA}(F, f) \Leftrightarrow \forall J \subseteq n \left(\left| \bigcup_{j \in J} F(j) \right| \geq |J| \right).$$

THEOREM 2. *Let $F = (F(i) : i \in n)$ be a finite family. Then*

$$\exists f \text{ IA}(F, f) \Leftrightarrow \neg \exists K \subseteq n \exists i \in n \setminus K \left(F(i) \subseteq \bigcup \text{rng } F \upharpoonright K \text{ \& } F \upharpoonright K \text{ critical} \right).$$

Proof. The one direction of Theorem 2 is Lemma 1. The other direction follows by an argument of the referee. Suppose F did not have an injective choice function. Then by P. Hall's theorem, there exists $K \subseteq n$ such that $|\bigcup_{i \in K} F(i)| < |K|$. Choose K minimal with respect to this property. Thus $|\bigcup \{F(i) : i \in K \setminus \{j\}\}| \geq |K \setminus \{j\}| = |K| - 1$ for all $j \in K$. Therefore $|K| - 1 \leq |\bigcup \{F(i) : i \in K \setminus \{j\}\}| \leq |\bigcup \{F(i) : i \in K\}| \leq |K| - 1$, so that $\bigcup \{F(i) : i \in K \setminus \{j\}\} = \bigcup \{F(i) : i \in K\}$ and $F(j) \subseteq \bigcup \{F(i) : i \in K \setminus \{j\}\}$ for all $j \in K$. But by minimality of K , $|\bigcup_{i \in I} F(i)| \geq |I|$ for all $I \subseteq K \setminus \{j\}$ so that $(F(i) : i \in K \setminus \{j\})$ is critical by Theorem 1 with $F(j) \subseteq \bigcup \{F(i) : i \in K \setminus \{j\}\}$ for all $j \in K$.

It is not difficult to prove:

LEMMA 2. *Let $F = (F(i) : i \in I)$ be a family which has an injective choice function (i.c.f.), and let \mathfrak{K} be a subset of $P(I)$. If $F \upharpoonright K$ is critical for all $K \in \mathfrak{K}$, then $(F(k) : k \in \bigcup \mathfrak{K})$ is a critical subfamily of F .*

LEMMA 3. *Let $F = (F(i) : i \in I)$ be a family, let $K \subseteq I$, $i_0 \in I \setminus K$, $F \upharpoonright K$ critical, $F(i_0) \subseteq \bigcup \text{rng } F \upharpoonright K$, and let $k_0 \in K$. Then*

$$\text{IA}(F \upharpoonright (K \setminus \{k_0\}) \cup \{i_0\}, f) \Rightarrow \text{rng } f = \bigcup \text{rng } F \upharpoonright K.$$

Proof. We use a zick-zack argument. Let f be an i.c.f. of

$$F \upharpoonright (K \setminus \{k_0\}) \cup \{i_0\}$$

and g an i.c.f. of $F \upharpoonright K$. We define a zick-zack (x_n) by induction. Let $x_0 = i_0$. If x_n has been defined, we let x_{n+1} be undefined, if $x_n = k_0$. The case $x_n \neq k_0$ implies $x_n \in \text{dom } f$, and since $F \upharpoonright K$ is critical, $f(x_n)$ is an element of $\text{rng } g$. Therefore we define $x_{n+1} = g^{-1}(f(x_n))$, if $x_n \neq k_0$. The zick-zack (x_n) is a finite sequence. Assume that (x_n) is not finite. Then $(g \setminus \{(x_n, g(x_n)) : 1 \leq n < \omega\}) \cup \{(x_n, f(x_n)) : n < \omega\}$ is an i.c.f. of $F \upharpoonright K \cup \{i_0\}$. Contradiction! Let $x_{n_0} = k_0$. The function

$$h = (f \setminus \{(x_n, f(x_n)) : 0 \leq n < n_0\}) \cup \{(x_n, g(x_n)) : 1 \leq n \leq n_0\}$$

is an i.c.f. of $F \upharpoonright K$ and $\text{rng } f = \text{rng } h = \bigcup \text{rng } F \upharpoonright K$.

COROLLARY. Let $F = (F(i): i \in I)$ be a family, $K \subseteq I$, $i_0 \in I \setminus K$, $F \upharpoonright K$ critical, $F(i_0) \subseteq \bigcup \text{rng } F \upharpoonright K$, and let $k_0 \in K$. If there is any i.c.f. of

$$F \upharpoonright (K \setminus \{k_0\}) \cup \{i_0\},$$

then $F \upharpoonright (K \setminus \{k_0\}) \cup \{i_0\}$ is a critical family.

We call a family with only finite members a *Hall-family*.

THEOREM 3 (M. Hall). A Hall-family F possesses an i.c.f. iff every finite subfamily of F possesses an i.c.f.

If $F = (F(i): i \in \omega)$ is a countable Hall-family, and if we define $f \leq g$ iff $f \subseteq g$ for $f, g \in \{h: \exists n \in \omega \text{ } IA(F \upharpoonright n, h)\}$, then M. Hall's result is an easy consequence of König's infinity lemma. You find a proof of Theorem 3 in the excellently written book of L. Mirsky.

THEOREM 4. Let $F = (F(i): i \in I)$ be a family, let $n \subseteq I$, and let $F \upharpoonright I \setminus n$ be a Hall-family. Then

$$\exists f \text{ } IA(F, f) \Leftrightarrow \neg \exists K \subseteq I \exists i \in I \setminus K (F(i) \subseteq \bigcup \text{rng } F \upharpoonright K \& F \upharpoonright K \text{ critical}).$$

Proof. By induction. For $n = 0$, the assertion follows from Theorems 2 and 3. Suppose that the assertion holds for n . Let $F = (F(i): i \in I)$ be a family, let $n + 1 \subseteq I$ and $F \upharpoonright I \setminus n + 1$ be a Hall-family, and let F satisfy the condition of Theorem 4. By the induction hypothesis, $F \upharpoonright I \setminus \{n\}$ has an i.c.f. Let us assume that F has no i.c.f. and let $x \in F(n)$.

Case 1. $\exists i \in I \setminus \{n\} F(i) = \{x\}$. Let i_0 be such an i . Then $H = \{(i_0, F(i_0))\}$ is a critical subfamily and $x \in \bigcup \text{rng } H$.

Case 2. $\forall i \in I \setminus \{n\} F(i) \neq \{x\}$. Let $F^*(i) = F(i) \setminus \{x\}$ for $i \in I \setminus \{n\}$. The family $F^* = (F^*(i): i \in I \setminus \{n\})$ has no i.c.f. By induction hypothesis, there exists an element $i_0 \in I \setminus \{n\}$ and a subset $K \subseteq I \setminus \{n\}$ such that $i_0 \notin K$, $F^* \upharpoonright K$ is critical and $F^*(i_0) \subseteq \bigcup \text{rng } F^* \upharpoonright K$. We shall prove that $(F(j): j \in K \cup \{i_0\})$ is critical. $F \upharpoonright K \cup \{i_0\}$ has an i.c.f. since

$$F \upharpoonright K \cup \{i_0\} \subseteq F \upharpoonright I \setminus \{n\}.$$

Assume that f is an i.c.f. of $F \upharpoonright K \cup \{i_0\}$, and let $y \in (\bigcup \text{rng } F \upharpoonright K \cup \{i_0\}) \setminus \text{rng } f$. It is $x \neq y$, since $F^* \upharpoonright K \cup \{i_0\}$ possesses no i.c.f. Assume $f(i_0) = x$. If $y \in F^*(i_0)$, then $F^* \upharpoonright K \cup \{i_0\}$ has an i.c.f. If $y \notin F^*(i_0)$, then $F^* \upharpoonright K$ is not critical. In every case there is a contradiction and therefore $f(i_0) \neq x$. Let $k_0 \in K$ and $f(k_0) = x$. Since $f^* = f \upharpoonright (K \setminus \{k_0\}) \cup \{i_0\}$ is an i.c.f. of

$$F^* \upharpoonright (K \setminus \{k_0\}) \cup \{i_0\},$$

it follows by Lemma 3 $\text{rng } f = \bigcup \text{rng } F^* \upharpoonright K$. $x \neq y$ implies

$$y \in \bigcup \text{rng } F^* \upharpoonright K \cup \{i_0\}.$$

Otherwise there is $\bigcup \text{rng } F^* \upharpoonright K \cup \{i_0\} = \text{rng } F^* \upharpoonright K$. Therefore y is an element of $\bigcup \text{rng } F^* \upharpoonright K = \text{rng } f^*$. Contradiction! We have proved that $F \upharpoonright K \cup \{i_0\}$ is a critical family. Hence x is an element of $\bigcup \text{rng } F \upharpoonright K \cup \{i_0\}$.

Each $x \in F(n)$ and each case leads to a critical subfamily $F \upharpoonright K_x$ of $F \upharpoonright I \setminus \{n\}$. By Lemma 2 $\bigcup \{F \upharpoonright K_x : x \in F(n)\}$ is a critical family and $F(n)$ is a subset of $\bigcup \text{rng } \bigcup \{F \upharpoonright K_x : x \in F(n)\}$, contrary to assumption. Hence F possesses an i.c.f.

Remark. This argument is also a proof of Theorem 2 without using Hall's theorem. Then Hall's theorem is a corollary of Theorem 2.

LEMMA 4. (Replacement Lemma). *Let $F = (F(i) : i \in I)$ be a family, let $K \subseteq I$, $i_0 \in I \setminus K$, $F(i_0) \subseteq \bigcup \text{rng } F \upharpoonright K$ and $F \upharpoonright K$ critical. Then there exists an element $k_0 \in K$ such that $F \upharpoonright (K \setminus \{k_0\}) \cup \{i_0\}$ is critical.*

Proof. Case 1. $\exists k \in K \exists f IA(F \upharpoonright (K \setminus \{k\}) \cup \{i_0\}, f)$. Then by the corollary of Lemma 3, there is nothing to prove. Case 2.

$$\neg \exists k \in K \exists f IA(F \upharpoonright (K \setminus \{k\}) \cup \{i_0\}, f).$$

Let $(k_\alpha)_{\alpha < \beta}$ be a bijection of the ordinal β onto the set K . Assume that there exists an element $\alpha < \beta$ and an i.c.f. of $F \upharpoonright K$ such that $f(k_\alpha)$ is an element of $F(i_0)$. Then $F \upharpoonright (K \setminus \{k_\alpha\}) \cup \{i_0\}$ possesses an i.c.f. Contradiction! Hence $\{f(k_\alpha) : IA(F \upharpoonright K, f)\} \cap F(i_0) = \emptyset$ for all $\alpha < \beta$. This means that

$$\bigcup_{\alpha < \beta} \{f(k_\alpha) : IA(F \upharpoonright K, f)\} \cap F(i_0) = \emptyset.$$

Since $\bigcup_{\alpha < \beta} \{f(k_\alpha) : IA(F \upharpoonright K, f)\} = \bigcup \text{rng } F \upharpoonright K$, we get a contradiction to $\emptyset \neq F(i_0) \subseteq \bigcup \text{rng } F \upharpoonright K$. We have proved that case 2 is impossible.

THEOREM 5. *Let $F = (F(i) : i \in I)$ be a family, let $n \subseteq I$, and let $F \upharpoonright I \setminus n$ be a Hall-family. Then $\exists f IA(F, f) \Leftrightarrow \bigcup \{G : G \subseteq F \text{ \& } G \text{ critical}\}$ critical.*

Proof. The direction " \Rightarrow " follows from Lemma 2. Suppose that F possesses no i.c.f. By Theorem 4 there is a set $K \subseteq I$ and an element $i_0 \in I \setminus K$ such that $F(i_0) \subseteq \bigcup \text{rng } F \upharpoonright K$ and $F \upharpoonright K$ is critical. By the replacement lemma, there exists an element $k_0 \in K$ such that $F \upharpoonright (K \setminus \{k_0\}) \cup \{i_0\}$ is a critical family. Since $(F \upharpoonright (K \setminus \{k_0\}) \cup \{i_0\}) \cup F \upharpoonright K = F \upharpoonright K \cup \{i_0\}$ possesses no i.c.f., the family $\bigcup \{G \subseteq F : G \text{ critical}\}$ is not critical.

LEMMA 5 (Brualdi, Scrimger, Folkman, and Steffens). *If the Hall-family F possesses an i.c.f., then $\bigcap \{\text{rng } f : IA(F, f)\} = \bigcup \{\text{rng } G : G \subseteq F \text{ critical \& } G \text{ finite}\}$.*

Proof. Since the inclusion " \supseteq " is obvious, let $x_0 \in \bigcap \{\text{rng } f : IA(F, f)\}$. If there is an element $i_0 \in I$ such that $F(i_0) = \{x_0\}$, then $x_0 \in \bigcap \{\text{rng } G : G \subseteq F \text{ critical \& } G \text{ finite}\}$. Let $F(i) \neq \{x_0\}$ for all $i \in I$. The family

$$F^* = (F(i) \setminus \{x_0\} : i \in I)$$

possesses no i.c.f., and therefore there exists, by Theorem 3, a finite subfamily $(F^*(j) : j \in J)$ which possesses no i.c.f. A Hall-argument (Theorem 1) shows that $F \upharpoonright J$ is a critical family and x_0 is an element of $\bigcup \text{rng } F \upharpoonright J$.

COROLLARY. *If F is a critical Hall-family, then $F = \bigcup \{G \subseteq F : G \text{ critical \& } G \text{ finite}\}$.*

Now there is no difficulty to extend the result of Lemma 5 to families with finitely many infinite members (without the finite property of G). As an application of Lemma 2, Theorem 4, Lemma 5, and the corollary to Lemma 5, we give a new proof of a theorem of J. Folkman.

THEOREM 6 (J. Folkman). *Let $F = (F(i) : i \in I)$ be a family, $n \subseteq I$, and $F \upharpoonright I \setminus n$ a Hall-family. Then $\exists f IA(F, f) \Leftrightarrow \forall K \forall J \forall r \in \omega (K \subseteq J \subseteq I \& \bigcup \text{rng } F \upharpoonright J \setminus \bigcup \text{rng } F \upharpoonright K \text{ finite \& } |J \setminus K| \geq r + |\bigcup \text{rng } F \upharpoonright J \setminus \bigcup \text{rng } F \upharpoonright K| \Rightarrow \exists S \in P_\omega(K) \forall T (S \subseteq T \subseteq K \Rightarrow |\bigcup \text{rng } F \upharpoonright T| \geq r + |T|))$.*

Proof. " \Rightarrow ", look at [2]. We prove the other direction of Theorem 6 by induction. Let $n = 0$ and suppose that F has no i.c.f. By Theorem 3 there is a finite subset L of I such that $F \upharpoonright L$ possesses no i.c.f. By Theorem 4 there is a set $K \subseteq L$, an element $1_0 \in L \setminus K$ such that $F(1_0) \subseteq \bigcup \text{rng } F \upharpoonright K$ and $|\bigcup \text{rng } F \upharpoonright K| = |K|$. If $J = K \cup \{1_0\}$, then

$$|J \setminus K| = 1 + |\bigcup \text{rng } F \upharpoonright J \setminus \bigcup \text{rng } F \upharpoonright K|.$$

Let S be a subset of K . Since $|\bigcup \text{rng } F \upharpoonright K| < 1 + |K|$, we choose for the set T the set K .

Now suppose that the assertion holds for every family with n many infinite sets. Let F be a family of $n + 1$ many infinite sets which satisfies the condition of Theorem 6. Assume that F possesses no i.c.f. By Lemma 2 $G = \bigcup \{H : H \subseteq F \upharpoonright I \setminus \{0\} \text{ critical}\}$ is a critical subfamily of F . Then $F(0) \subseteq \bigcap \{\text{rng } f : IA(F \upharpoonright I \setminus \{0\}, f)\} = \bigcup \{\text{rng } H : H \subseteq F \upharpoonright I \setminus \{0\} \text{ critical}\} = \bigcup \text{rng } G$. We note that $F(i) \in \text{rng } G$ for all $i \in \{1, \dots, n\}$, otherwise there is a

contradiction by the induction hypothesis. Let f be an i.c.f. of G and define for $i \in \text{dom } G \setminus \{1, \dots, n\}$ $G^*(i) = F(i) \setminus \{f(j) : j \in \{1, \dots, n\}\}$. The family G^* is a critical family of finite sets, since G is critical. By the corollary to Lemma 5, G^* is the union of all finite, critical subfamilies of G^* . Hence there exists to every finite subfamily H^* of G^* a finite, critical family K^* such that $H^* \subseteq K^* \subseteq G^*$. Let

$$r = \left| \{f(i) : i \in \{1, \dots, n\}\} \cap \bigcup \text{rng } F \upharpoonright \text{dom } G^* \right|,$$

$$J = \text{dom } G \cup \{0\}, \quad \text{and} \quad K = \text{dom } G^*.$$

$|\bigcup \text{rng } F \upharpoonright J \setminus \bigcup \text{rng } F \upharpoonright K| = n - r$ implies

$$|J \setminus K| = n + 1 = (r + 1) + \left| \bigcup \text{rng } F \upharpoonright J \setminus \bigcup \text{rng } F \upharpoonright K \right|.$$

Let $S \in P_\omega(K)$. There exists a finite, critical subfamily H^* of G^* such that $G^* \upharpoonright S \subseteq H^*$. Let $T = \text{dom } H^*$, then $S \subseteq T \subseteq K$ and

$$|\bigcup \text{rng } F \upharpoonright T| < (r + 1) + |T|,$$

contrary to assumption.

We now present a generalization of a theorem of H. A. Jung and R. Rado.

THEOREM 7. *Let $F = (F(i) : i \in I)$ be a family, let $I_0 \subseteq I$, and let $F \upharpoonright I \setminus I_0$ be a Hall-family. Then*

$$\begin{aligned} \exists g \text{ } IA(F, g) &\Leftrightarrow \exists f \left(IA(F \upharpoonright I_0, f) \ \& \ \forall J \in P_\omega(I \setminus I_0) \right. \\ &\quad \times \left. \left(\left| \bigcup \{F(j) : j \in J\} \cap \text{rng } f \right| = k \Rightarrow \left| \bigcup_{j \in J} F(j) \right| \geq |J| + k \right) \right). \end{aligned}$$

Proof. “ \Rightarrow ”. Let g be an i.c.f. of F and $f = g \upharpoonright I_0$. Then f is an i.c.f. of $F \upharpoonright I_0$. If there is a finite subset J of I such that $k = |\bigcup_{j \in J} F(j) \cap \text{rng } f|$ and $|\bigcup \{F(j) : j \in J\}| < |J| + k$, then f and therefore g are not injective functions. Contradiction!

“ \Leftarrow ”. Let F be a family which satisfies the condition of Theorem 7, let f be an i.c.f. of $F \upharpoonright I_0$, and let for $i \in I \setminus I_0$ $F^*(i) = F(i) \setminus \text{rng } f$. If $J \in P_\omega(I \setminus I_0)$, then $|\bigcup_{j \in J} F^*(j)| \geq |J|$. By P. Hall's theorem and by Theorem 3, F^* possesses an i.c.f. Hence there exists an i.c.f. of the family F .

COROLLARY (Jung-Rado). *Let $F = (F(i) : i \in I^*)$ be a family, $i_0 \in I^*$, $I = I^* \setminus \{i_0\}$, and let $F \upharpoonright I$ be a Hall-family. Then $\exists f \text{ } IA(F, f) \Leftrightarrow \forall J \in P_\omega(I)$ $(|\bigcup \text{rng } F \upharpoonright J| \geq |J|) \ \& \ F(i_0) \not\subseteq \bigcup \{\text{rng } G : G \subseteq F \text{ critical} \ \& \ G \text{ finite}\}$.*

Proof. We say that “ F satisfies \mathcal{H} ” if it is valid:

$$\forall J \in P_\omega(I) (|\bigcup \text{rng } F \upharpoonright J| \geq |J|).$$

$\exists f \text{ IA}(F, f)$

$$\Leftrightarrow \exists x \in F(i_0) \forall J \in P_\omega(I)$$

$$\times \left(\left| \bigcup \{F(j): j \in J\} \cap \{x\} \right| = k \rightarrow \left| \bigcup \{F(j): j \in J\} \right| \geq |J| + k \right)$$

$$\Leftrightarrow F \text{ satisfies } \mathcal{H} \text{ \& } \exists x \in F(i_0) \forall J \in P_\omega(I)$$

$$\times \left(x \in \bigcup \text{rng } F \upharpoonright J \Rightarrow \left| \bigcup \text{rng } F \upharpoonright J \right| \geq |J| + 1 \right)$$

$$\Leftrightarrow F \text{ satisfies } \mathcal{H} \text{ \& } \exists x \in F(i_0) \neg \exists J \in P_\omega(I)$$

$$\times \left(x \in \bigcup \text{rng } F \upharpoonright J \text{ \& } \left| \bigcup \text{rng } F \upharpoonright J \right| = |J| \right)$$

$$\Leftrightarrow F \text{ satisfies } \mathcal{H} \text{ \& } F(i_0) \not\subseteq \bigcup \{\text{rng } F \upharpoonright J: J \in P_\omega(I) \text{ \& } F \upharpoonright J \text{ critical}\}$$

$$\Leftrightarrow F \text{ satisfies } \mathcal{H} \text{ \& } F(i_0) \not\subseteq \bigcup \{\text{rng } G: G \subseteq F \text{ critical \& } G \text{ finite}\}.$$

Remark. (1) We do not know any criterion for families with only infinite members.

(2) The family $F = \{(\alpha, \omega): \alpha < \omega_1\}$ possesses neither a nonempty, critical subfamily nor an injective choice function. But we do not know if Theorem 4 can be extended to families with countably many infinite members. You will find further problems in [7].

Note Added in Proof. Recently question (2) was positively answered by K.-P. Podewski and the author.

REFERENCES

1. R. A. BRUALDI AND E. B. SCRIMGER, Exchange systems, matchings and transversals, *J. Combinatorial Theory* **5** (1968), 244–257.
2. J. FOLKMAN, Transversals of infinite families with finitely many infinite members, *J. Combinatorial Theory* **9** (1970), 200–220.
3. M. HALL, JR., Distinct representatives of subsets, *Bull. Amer. Math. Soc.* **54** (1948), 922–926.
4. P. HALL, On representatives of subsets, *J. Lond. Math. Soc.* **10** (1935), 26–30.
5. L. MIRSKY, “Transversal Theory,” Academic Press, New York, 1971.
6. R. RADO, Note on the transfinite case of Hall’s theorem on representatives, *J. Lond. Math. Soc.* **42** (1967), 321–324.
7. K. STEFFENS, *Injektive Auswahlfunktionen*, Schriften aus dem Gebiet der Angewandten Mathematik, Aachen 1972.